



Acciarresi, M. et al. (2017) Prestroke CHA2DS2-VASc score and severity of acute stroke in patients with atrial fibrillation: findings from RAF study. *Journal of Stroke and Cerebrovascular Diseases*, 26(6), pp. 1363-1368. (10.1016/j.jstrokecerebrovasdis.2017.02.011)

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/138063/>

Deposited on: 15 September 2017

Enlighten – Research publications by members of the University of Glasgow
<http://eprints.gla.ac.uk>

Pre-stroke CHA₂DS₂-VASc score and severity of acute stroke in patients with atrial fibrillation: findings from RAF study

Monica Acciarresi MD¹, Maurizio Paciaroni MD¹, Giancarlo Agnelli MD¹, Nicola Falocci, PhD¹, Valeria Caso MD, PhD¹, Cecilia Becattini MD¹, Simona Marcheselli MD², Christina Rueckert MD³, Alessandro Pezzini MD⁴, Andrea Morotti MD⁴, Paolo Costa MD⁴, Alessandro Padovani MD, PhD⁴, Laszló Csiba MD⁵, Lilla Szabó MD⁵, Sung-Il Sohn MD, PhD⁶, Tiziana Tassinari MD⁷, Azmil H Abdul-Rahim MD⁸, Patrik Michel, PD-MER⁹, Maria Cordier MD⁹, Peter Vanacker MD¹⁰, Suzette Remillard MD⁹, Andrea Alberti MD¹, Michele Venti MD, PhD¹, Cataldo D'Amore MD¹, Umberto Scoditti MD¹¹, Licia Denti MD¹², Giovanni Orlandi MD¹³, Alberto Chiti MD¹³, Gino Gialdini MD¹³, Paolo Bovi MD¹⁴, Monica Carletti MD¹⁴, Alberto Rigatelli MD¹⁴, Jukka Putaala MD¹⁵, Turgut Tatlisumak MD^{15,16}, Luca Masotti MD¹⁷, Gianni Lorenzini MD¹⁷, Rossana Tassi MD¹⁸, Francesca Guideri MD¹⁸, Giuseppe Martini MD¹⁸, Georgios Tsivgoulis MD^{19,20,21}, Kostantinos Vadikolias MD¹⁹, Chrissoula Liantinioti MD²¹, Francesco Corea MD, PhD²², Massimo Del Sette MD²³, Walter Ageno MD²⁴, Maria Luisa De Lodovici MD²⁵, Giorgio Bono MD²⁵, Antonio Baldi MD²⁶, Sebastiano D'Anna MD²⁶, Simona Sacco MD²⁷, Antonio Carolei²⁷, Cindy Tiseo MD²⁷, Davide Imberti MD²⁸, Dorjan Zabzuni MD²⁸, Boris Doronin MD²⁹, Vera Volodina MD²⁹, Domenico Consoli MD³⁰, Franco Galati MD³⁰, Alessio Pieroni MD³¹, Danilo Toni MD, PhD³¹, Serena Monaco MD³², Mario Maimone Baronello MD³², Kristian Barlinn MD³³, Lars-Peder Pallesen MD³³, Jessica Kepplinger MD³³, Ulf Bodechtel MD³³, Johannes Gerber MD³³, Dirk Deleu, MD³⁴, Gayane Melikyan MD³⁴, Faisal Ibrahim MD³⁴, Naveed Akhtar MD³⁴, Maria Giulia Mosconi MD¹, Kennedy R Lees MD⁸

¹Stroke Unit and Division of Cardiovascular Medicine, University of Perugia, Italy

²Neurologia d'urgenza e Stroke Unit, Istituto Clinico Humanitas, Rozzano, Milano, Italy

³Abteilung für Neurologie, Oberschwabenklinik gGmbH, Ravensburg, Germany

⁴Department of Clinical and Experimental Sciences, Neurology Unit, University "Health and Wealth" of Brescia, Italy

⁵Stroke Unit, University of Debrecen, Hungary

⁶Department of Neurology, Keimyung University School of Medicine, Daegu, South Korea

⁷Stroke Unit-Department of Neurology, Santa Corona Hospital, Pietra Ligure (Savona), Italy

⁸Medical School and Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, United Kingdom.

⁹Centre Cérébrovasculaire, Service de Neurologie, Département des Neurosciences Cliniques Centre Hospitalier Universitaire Vaudois, Lausanne (Switzerland)

- ¹⁰Department of Neurology, Born Bunge Institute, Antwerp University Hospital, Antwerp, Belgium
¹¹Stroke Unit, Neuroscience Department, University of Parma, Italy
¹²Stroke Unit - Dipartimento Geriatrico Riabilitativo – University of Parma, Italy
¹³Clinica Neurologica – Azienda Ospedaliero-Universitaria, Pisa, Italy
¹⁴SSO Stroke Unit, UO Neurologia, DAI di Neuroscienze, AOUI Verona, Italy
¹⁵Department of Neurology, Helsinki University Central Hospital, Helsinki, Finland
¹⁶Institute of Neuroscience and Physiology, Sahlgrenska Academy at University of Gothenburg and Department of Neurology, Sahlgrenska University Hospital, Gothenburg, Sweden
¹⁷Department of Internal Medicine, Cecina Hospital, Cecina, Livorno, Italy
¹⁸Stroke Unit, AOU Senese, Siena, Italy
¹⁹Department of Neurology, Democritus University of Thrace, University Hospital of Alexandroupolis, Greece
²⁰International Clinic Research Center, St. Anne's University Hospital Brno, Brno, Czech Republic
²¹Second Department of Neurology, "Attikon" Hospital, University of Athens, School of Medicine, Athens, Greece
²²UO Gravi Cerebrolesioni, San Giovanni Battista Hospital, Foligno
²³Stroke Unit, Department of Neurology, Sant'Andrea Hospital, La Spezia, Italy
²⁴Department of Internal Medicine, Insubria University, Varese, Italy
²⁵Stroke Unit, Neurology, Insubria University, Varese, Italy
²⁶Stroke Unit, Ospedale di Portogruaro, Portogruaro (Venice), Italy
²⁷Department of Neurology, University of L'Aquila, Italy
²⁸Department of Internal Medicine, Ospedale Civile di Piacenza, Italy
²⁹Municipal Budgetary Healthcare Institution of Novosibirsk. City Clinical Hospital #1. Novosibirsk (Russia)
³⁰Stroke Unit, Jazzolino Hospital, Vibo Valentia, Italy
³¹Department of Neurology and Psychiatry, Sapienza University of Rome, Italy
³²Stroke Unit, Ospedale Civico, Palermo
³³Department of Neurology, Dresden University Stroke Center, Dresden, Germany
³⁴Neurology, Hamad Medical Corporation, Doha, Qatar

Corresponding author:

Monica Acciarresi

Stroke Unit and Division of Internal and Cardiovascular Medicine

University of Perugia, Santa Maria della Misericordia Hospital

Via G. Dottori 1, Perugia

06100 – Italy

Email: macun77@hotmail.com

Tel and fax: ++39(0)75.5782765

Key words: ischemic stroke, CHA2DS2-VASc score, severity, outcome, atrial fibrillation, scores

Word count: 1.808

Tables: 2

Figures: 2

Abstract

Background and Purpose: The aim of this study was to investigate for a possible association between both pre-stroke CHA₂DS₂-VASc score and the severity of stroke at presentation as well as disability and mortality at 90 days in patients with acute stroke and atrial fibrillation (AF).

Methods: This prospective study enrolled consecutive patients with acute ischemic stroke, AF and assessment of pre-stroke CHA₂DS₂-VASc score. Severity of stroke was assessed at admission by the National Institute of Health Stroke Scale (NIHSS) score, (severe stroke: NIHSS \geq 10). Disability and mortality at 90 days were assessed by the modified Rankin Scale (mRS $<$ 3 or \geq 3). Multiple logistic regression was used to correlate pre-stroke CHA₂DS₂-VASc and severity of stroke as well as disability and mortality at 90 days.

Results: Of the 1.020 patients included in the analysis, 606 patients had an admission NIHSS score lower and 414 patients higher than 10. At 90 days, 510 patients had mRS \geq 3. A linear correlation was found between pre-stroke CHA₂DS₂-VASc score and severity of stroke (p=0.001). On multivariate analysis, CHA₂DS₂-VASc score correlated with the severity of stroke (p=0.041) and adverse functional outcome (mRS \geq 3) (p=0.001). A logistic regression with the ROC graph procedure (C statistic) evidenced an area under the curve of 0.60 (p=0.0001) for severe stroke. Furthermore, a correlation was found between pre-stroke CHA₂DS₂-VASc score and the lesion size.

Conclusions: In patients with AF, in addition to the risk of stroke, a high CHA₂DS₂-VASc score was independently associated with both stroke severity at onset and disability and mortality at 90 days.

Background and Purpose

In patients with atrial fibrillation (AF), current guidelines recommend using the CHA₂DS₂-VASc score to assess the risk of stroke¹. In retrospective studies CHA₂DS₂-VASc score have been suggested as a predictor of severity of stroke at admission and poor outcome²⁻⁴.

The aim of this study was to investigate for a possible association between both pre-stroke CHA₂DS₂-VASc score and the severity of stroke at presentation as well as disability and mortality at 90 days in patients with acute stroke and atrial fibrillation (AF).

Methods

Data for this analysis were extracted from the database of a prospective multicentre study which had enrolled consecutive patients with acute stroke and AF (the RAF study)⁵. This study, carried out between January 2012 and March 2014, enrolled 1,029 consecutive patients from 29 Stroke Units throughout Europe and Asia.

Pre-stroke CHA₂DS₂-VASc score was evaluated as previously described⁵. On admission, the severity of acute stroke was assessed using the National Institutes of Health Stroke Scale (NIHSS); all investigators were certified on the use of this scale. Disability and mortality at 90 days were assessed using the modified Rankin Scale (mRS).

Data on known stroke risk factors were collected as reported in the main paper⁵.

A cerebral computed tomography (CT) or magnetic resonance (MR) was performed on admission for all patients to exclude intracranial hemorrhage. A second cerebral CT scan or MR was performed 48-72 h from stroke onset. The sites and sizes of the qualifying infarcts were determined based on standard templates ^{6,7} as previously described ⁵.

Statistical analysis

The primary pre-specified a priori study question was if CHA₂DS₂-VASc score had been associated with stroke severity assessed by NIHSS at onset. Then, as a post-hoc analysis,

possible correlations between CHA₂DS₂-VASc scores and outcome assessed by mRS and between CHA₂DS₂-VASc scores and lesion size were investigated for.

The admission NIHSS score was evaluated both as a continuous variable for the correlation coefficient r analysis, and dichotomized variables (severe stroke NIHSS \geq 10) for the multivariate analysis.

The correlation coefficient r , Pearson product-moment correlation coefficient, was used to measure the strength of the linear association between CHA₂DS₂-VASc before the event and NIHSS score on admission (as continuous variable).

Correlations between pre-stroke CHA₂DS₂-VASc and severity of stroke were sought by multiple logistic regression after adjusting for the following variables: smoking, hyperlipidemia, alcohol abuse and use of statins in addition to CHA₂DS₂-VASc.

Thereafter, the probability of a receiver-operating characteristic (ROC) curve against NIHSS \geq 10 as dependent variable was plotted. The area under this curve suggests an ability of the CHA₂DS₂-VASc score to predict for severe stroke, which is also referred to as the C-statistic (Harrell's C).

In addition, a multiple logistic regression model including components singularly of the CHA₂DS₂-VASc was performed.

Stroke was defined as either non-disabling (mRS 0 to 2) or disabling (mRS 3 to 5).

The factors evaluated as independent predictors of 3 month adverse outcome (defined as modified Rankin scale \geq 3 or death) were assessed using multiple logistic regression analysis. The variables included in the model were CHA₂DS₂-VASc score, cardiovascular risk factors, reperfusion therapy and severity of stroke on admission according to NIHSS score.

The sites and sizes of the qualifying infarcts showed at CT scan were determined based on standard templates as follows: (1) small, when a lesion was \leq 1.5 cm in the anterior or posterior circulation, (2) medium, when a lesion was in a cortical superficial branch of middle cerebral artery [MCA], in the MCA deep branch, in the internal border zone territories, in a cortical superficial branch of posterior cerebral artery [PCA], in the PCA

branch or in a cortical superficial branch of the anterior cerebral artery [ACA]), (3) large anterior, when a lesion involved the complete territory of ACM, ACP, or ACA, in 2 cortical superficial branches of MCA, in a cortical superficial branch of MCA associated to the MCA deep branch, or in more than 1 artery territory [eg, MCA associated to ACA territories]], (4) large posterior, when a lesion was ≥ 1.5 cm in the brain stem or cerebellum⁸.

A possible correlation between lesion size and CHA₂DS₂-VASc score was investigated for using a χ -squared (p for trend).

Results:

Overall, 1,029 consecutive patients were included in the study (mean age 77.2 ± 9.5 years; 560 females and 469 males). The distribution of CHA₂DS₂-VASc score in the study patients is reported in the Table 1.

Nine patients were excluded from the analysis as their NIHSS score was not available. Of the 1,020 patients included in the analysis, 606 patients had an admission NIHSS score lower and 414 patients higher than 10. The mean NIHSS score for each of the CHA₂DS₂-VASc scores are reported in the Table 1 ($p=0.071$). A linear correlation was found between severity of stroke and CHA₂DS₂-VASc score (r^2 0.010, $p=0.001$) (fig. 1). On multivariate analysis, CHA₂DS₂-VASc score correlated with the severity of stroke (OR 1.084, 95% CI 1.003-1.172, $p=0.041$, for each one point increase). Considering the modest correlation, a logistic regression with a ROC graph procedure, to get the c statistic, was performed and it evidenced that the area under the curve was 0.60 (0.56-0.63), $p=0.0001$ for severe stroke.

When factors included in the CHA₂DS₂-VASc score were assessed individually for a correlation between stroke severity, only increasing age and female sex resulted being significantly correlated to stroke severity according to NIHSS ($p=0.007$ and $p=0.001$, respectively) (Table 2).

A correlation was also found between pre-stroke CHA₂DS₂-VASc score and lesion size. After excluding patients treated with revascularization, 70% of patients with CHA₂DS₂-

VASc score of 8 or 9 had a medium-large lesion, while 70% of patients with CHA₂DS₂-VASc score of 0 had a small lesion (Figure 2, p for trend=0.042).

At 90-days, 1,019 patients were available for the functional outcome analysis (10 patients were lost at follow-up). The correlation between pre-stroke CHA₂DS₂-VASc score and 90 day outcome is reported in Table 1.

On multivariate analysis, both NIHSS score on admission and CHA₂DS₂-VASc score were correlated with disability and mortality at 90 days (mRS \geq 3) (OR 1.236, 95% CI 1.197-1.277, p<0.0001; OR 1.278; 95% CI 1.100-1.484, p=0.001 respectively for each one point increase).

The administration of thrombolytic therapy and the use of statins at admission resulted being independently correlated with better outcome (OR 0.313, 95% CI 0.204-0.481, p<0.0001; OR 0.585, 95% CI 0.354-0.967, p=0.036; respectively).

Discussion:

The results of our study found correlations between pre-stroke CHA₂DS₂-VASc score and severity of stroke at presentation, according to NIHSS, as well as between CHA₂DS₂-VASc score and 90-day mortality and disability, as measured by mRS.

Therein, indicating that a higher CHA₂DS₂-VASc score is predictive of a worse outcome. Moreover, in post-hoc analyses we also found, first, a correlation between higher CHA₂DS₂-VASc score and greater lesion size. This finding could support previous studies that have reported an association between CHA₂DS₂-VASc score and major vessel occlusion in patients with acute ischemic stroke and AF⁹. Second, our study results also is in line with past studies reporting that among risk factors, age and female sex are more predictive of severe stroke^{10,11}.

A strength of this study was that several stroke assessment scores were prospectively utilized, and each of these scores were correlated to the CHA₂DS₂-VASc score. Whereas, several past studies were retrospective and used only a single assessment score^{2-4,12-14}.

In conclusion, we found an increasing CHA2DS2-VASc score in AF patients predicted a more severe stroke at presentation, leading to a higher rate of disability and mortality at 90 days. This result highlights the need for physicians to more regularly investigate for AF in patients with other known vascular risk factors, especially increasing age and female sex.

Table 1: Stroke severity (NIHSS) and 90 day outcome according to pre-stroke CHA₂DS₂-VASc score

CHA ₂ DS ₂ -VASc score	Pre-stroke CHA ₂ DS ₂ -VASc (n=1029)	NIHSS at admission (mean) [^] (n=1020 ^{**})	90-day mRS score≥3* (n=1019 ^{***})
0	17 (1.7%)	5.18 ± 5.75	4/17 (23.5%)
1	54 (5.2%)	8.30 ± 8.16	16/53 (30.2%)
2	91 (8.9%)	8.31 ± 6.70	34/90 (37.7%)
3	200 (19.4%)	8.86 ± 7.10	81/198 (40.9%)
4	243 (23.6%)	9.12 ± 7.22	125/242 (51.6%)
5	206 (20.0%)	9.67 ± 7.73	113/202 (55.9%)
6	129 (12.8%)	9.70 ± 6.50	78/129 (60.5%)
7	66 (6.4%)	10.30 ± 8.20	43/65 (66.1%)
8-9	23 (2.2%)	12.09 ± 6.68	16/23 (69.6%)

[^]p for trend=0.071

*p for trend=0.0001

^{**} Nine patients were excluded from the analysis as their NIHSS score was not available

^{***}1019 patients were available for the final functional outcome analysis (10 patients were lost at follow-up)

Table 2. Multivariate analysis for correlation of stroke severity and any of each component of CHA₂DS₂-VASC score.

	OR	P	95% CI
Age (for each year increase)	1.020	0.007	1.005-1.035
Sex	0.367	0.001	0.515-0.175
Diabetes	1.172	0.296	0.870-1.575
Hypertension	0.759	0.110	0.542-1.065
Stroke/TIA/Thromboembolism	0.972	0.849	0.723-1.306

History			
Congestive Heart Failure	1.049	0.777	0.753-1.462
Vascular Disease	1.034	0.832	0.760-1.406

Fig. 1. Linear correlation between NIHSS at admission and CHA2DS2-VASC score.





Figure 2: CHA₂DS₂-VASC score and lesion size (excluding patients treated with revascularization).



* p for trend=0.042

Disclosures

On behalf of all authors, the corresponding author states that there is no conflict of interest.

M. Paciaroni received honoraria as a member of the speaker bureau of Sanofi-Aventis, Boehringer Ingelheim, Bayer and Pfizer. G. Agnelli received honoraria as a member of the speaker bureau of Boehringer Ingelheim and Bayer. C. Becattini received honoraria as a member of the speaker bureau of Bristol Meyer Squibb and Bayer. P. Michel received Research Grant by Swiss National Science Foundation and Swiss Heart Foundation; he received speaker fees by Bayer, Boehringer Ingelheim, Covidien, St. Jude Medical; he received honoraria as advisory relationship by Pierre-Fabre, Bayer, Bristol Meyer Squibb, Amgen, and Boehringer Ingelheim. J. Putaala received honoraria for lectures related to atrial fibrillation and anticoagulants for Orion Pharma, Bristol Meyer Squibb, Pfizer,

Bayer, and Boehringer Ingelheim. T. Tatlisumak received honoraria as consultant or advisory relationship by Lundbeck and Boehringer Ingelheim. G. Tsivgoulis had research support by European Regional Development Fund, Project St. Anne's University Hospital, Brno, International Clinical Research Center (FNUSA-ICRC) (No. CZ.1.05/1.1.00/02.0123). D. Toni received honoraria as a member of speaker bureau and as advisory board of Boehringer Ingelheim and Bayer. The other authors report no conflicts.

References:

1. Camm a J, Lip GYH, De Caterina R, et al. 2012 focused update of the ESC Guidelines for the management of atrial fibrillation: an update of the 2010 ESC Guidelines for the management of atrial fibrillation--developed with the special contribution of the European Heart Rhythm Association. *Europace* 2012;14:1385–413.
2. Hong HJ, Kim YD, Cha M-J, et al. Early neurological outcomes according to CHADS2 score in stroke patients with non-valvular atrial fibrillation. *Eur J Neurol* 2012;19(2):284–90.
3. Giralt-Steinhauer E, Cuadrado-Godia E, Ois Á, et al. CHA(2)DS(2)-VASc score and prognosis in ischemic strokes with atrial fibrillation. *J Neurol* 2012;259(4):745–51.
4. Deguchi I, Hayashi T, Ohe Y, et al. The CHA(2)DS(2)-VASc score reflects clinical outcomes in nonvalvular atrial fibrillation patients with an initial cardioembolic stroke. *J Stroke Cerebrovasc Dis* 2013;22(8):e343–6.
5. Paciaroni M, Agnelli G, Falocci N, et al. Early Recurrence and Cerebral Bleeding in Patients With Acute Ischemic Stroke and Atrial Fibrillation: Effect of Anticoagulation and Its Timing: The RAF Study. *Stroke* 2015;46(8):2175–82.
6. Tatu L, Moulin T, Bogousslavsky J, Duvernoy H. Arterial territories of human brain: brainstem and cerebellum. *Neurology* 1996;47(5):1125–35.
7. Tatu L, Moulin T, Bogousslavsky J, Duvernoy H. Arterial territories of the human brain: cerebral hemispheres. *Neurology* 1998;50(6):1699–708.
8. Paciaroni M, Agnelli G, Corea F, et al. Early hemorrhagic transformation of brain infarction: rate, predictive factors, and influence on clinical outcome: results of a prospective multicenter study. *Stroke* 2008;39(8):2249–56.
9. Sakamoto Y, Sato S, Hama Y, Nagatsuka K, Minematsu K, Toyoda K. The relationship between the pre-admission CHA2DS2-VASc score and proximal artery occlusion in patients with acute stroke and atrial fibrillation. *Eur J Neurol* 2015;22(7):1081–7.
10. Bushnell CD, Lee J, Duncan PW, Newby LK, Goldstein LB. Impact of comorbidities on ischemic stroke outcomes in women. *Stroke* 2008;39(7):2138–40.
11. Santalucia P, Pezzella FR, Sessa M, et al. Sex differences in clinical presentation, severity and outcome of stroke: results from a hospital-based registry. *Eur J Intern Med* 2013;24(2):167–71.
12. Tu HTH, Campbell BC V, Meretoja A, et al. Pre-stroke CHADS2 and CHA2DS2-VASc scores are useful in stratifying three-month outcomes in patients with and without atrial fibrillation. *Cerebrovasc Dis* 2013;36(4):273–80.
13. Yaghi S, Sherzai A, Pilot M, Sherzai D, Elkind MS V. The CHADS2 Components Are Associated with Stroke-Related In-hospital Mortality in Patients with Atrial Fibrillation. *J Stroke Cerebrovasc Dis* 2015;24(10):2404–7.
14. Tanaka K, Yamada T, Torii T, et al. Pre-admission CHADS2, CHA2DS2-VASc, and R2CHADS2 Scores on Severity and Functional Outcome in Acute Ischemic Stroke with Atrial Fibrillation. *J Stroke Cerebrovasc Dis* 2015;24(7):1629–35.